

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Docket No.: C 2

Claims

Claim 1: A composition comprising:

(a) an oily substance; and

(b) a solubilizer containing:

(i) a first surfactant selected from the group consisting of a fatty alcohol ethoxylate corresponding to formula (I):

$$R^1-(C_2H_4O)_n-H$$
 wherein R^1 is a linear or branched, saturated or unsaturated alkyl moiety having from about 6 to 22 carbon atoms, and n is an integer from 1 to about 10, a fatty alcohol alkoxylate corresponding to formula (II):
$$R^2-(C_2H_4O)_n-(C_3H_6O)_m-H$$
 wherein R^2 is a linear or branched, saturated or unsaturated alkyl moiety having from about 6 to 22 carbon atoms, and n and m are, independently from one another, an integer from 1 to about 10, and mixtures thereof;

(ii) an alkyl and/or alkenyl oligoglycoside corresponding to formula (III):

$$R^3O-[G]_p$$
 wherein R^3 is a branched and unbranched alkyl and/or alkenyl radical having from about 4 to 22 carbon atoms, G is a sugar radical having 5 or 6 carbon atoms, and p is a number from 1 to 10;

(iii) a fatty alcohol ether sulfate corresponding to formula (IV):

$$R^4-(C_2H_4O)_n-SO_3K^+$$
 wherein R^4 is a linear or branched, saturated or unsaturated alkyl moiety having from about 6 to 22 carbon atoms, n is a number from 1 to about 10, and K^+ represents a cation; and

(iv) optionally, an acid component.

Claim 2: The composition of claim 1 wherein (i) is present in the solubilizer in an amount of from about 1 to 20% by weight, based on the weight of the solubilizer.

Claim 3: The composition of claim 1 wherein (i) is present in the solubilizer in an amount of from about 2 to 15% by weight, based on the weight of the solubilizer.

Claim 4: The composition of claim 1 wherein (i) is present in the solubilizer in an amount of from about 5 to 10% by weight, based on the weight of the solubilizer.

Docket No.: C2

Claim 5: The composition of claim 1 wherein (ii) is present in the solubilizer in an amount of from about 10 to 60% by weight, based on the weight of the solubilizer.

Claim 6: The composition of claim 1 wherein (ii) is present in the solubilizer in an amount of from about 25 to 50% by weight, based on the weight of the solubilizer.

Claim 7: The composition of claim 1 wherein (iii) is present in the solubilizer in an amount of from about 10 to 80% by weight, based on the weight of the solubilizer.

Claim 8: The composition of claim 1 wherein (iii) is present in the solubilizer in an amount of from about 25 to 60% by weight, based on the weight of the solubilizer.

Claim 9: The composition of claim 1 wherein (iii) is present in the solubilizer in an amount of from about 30 to 55% by weight, based on the weight of the solubilizer.

Claim 10: The composition of claim 1 wherein (iv) is present in the solubilizer in an amount of from about 0.1 to 3% by weight, based on the weight of the solubilizer.

Claim 11: The composition of claim 1 wherein the solubilizer is present in the composition in an amount of from about 1 to 30% by weight, based on the weight of the composition.

Claim 12: A process for solubilizing an oily substance comprising:

(a) providing an oily substance;

(b) providing a solubilizer containing:

(i) a first surfactant selected from the group consisting of a fatty alcohol ethoxylate corresponding to formula (I):

$R^1-(C_2H_4O)_n-H$, wherein R^1 is a linear or branched, saturated or unsaturated alkyl moiety having from about 6 to 22 carbon atoms, and n is an integer from 1 to about 10, a fatty alcohol alkoxyate corresponding to formula (II);

$R^2-(C_2H_4O)_n-(C_3H_6O)_m-H$, wherein R^2 is a linear or branched, saturated or unsaturated alkyl moiety having from about 6 to 22 carbon atoms,

Docket No.: C 2

and n and m are, independently from one another, an integer from 1 to about 10, and mixtures thereof;

(ii) an alkyl and/or alkenyl oligoglycoside corresponding to formula (III):

$R^3O-[G]_p$, wherein R^3 is a branched and unbranched alkyl and/or alkenyl radical having from about 4 to 22 carbon atoms, G is a sugar radical having 5 or 6 carbon atoms, and p is a number from 1 to 10;

(iii) a fatty alcohol ether sulfate corresponding to formula (IV):

$R^4-(C_2H_4O)_n-SO_3^-K^+$ wherein R^4 is a linear or branched, saturated or unsaturated alkyl moiety having from about 6 to 22 carbon atoms, n is a number from 1 to about 10, and K^+ represents a cation; and

(iv) optionally, an acid component; and

(c) combining (a) and (b).

Claim 13: The process of claim 12 wherein (i) is present in the solubilizer in an amount of from about 1 to 20% by weight, based on the weight of the solubilizer.

Claim 14: The process of claim 12 wherein (i) is present in the solubilizer in an amount of from about 5 to 10% by weight, based on the weight of the solubilizer.

Claim 15: The process of claim 12 wherein (ii) is present in the solubilizer in an amount of from about 10 to 60% by weight, based on the weight of the solubilizer.

Claim 16: The process of claim 12 wherein (ii) is present in the solubilizer in an amount of from about 25 to 50% by weight, based on the weight of the solubilizer.

Claim 17: The process of claim 12 wherein (iii) is present in the solubilizer in an amount of from about 10 to 80% by weight, based on the weight of the solubilizer.

Claim 18: The process of claim 12 wherein (iii) is present in the solubilizer in an amount of from about 30 to 55% by weight, based on the weight of the solubilizer.

Claim 19: The process of claim 12 wherein (iv) is present in the solubilizer in an amount of from about 0.1 to 3% by weight, based on the weight of the solubilizer.

Docket No.: C 2

Claim 20: The process of claim 12 wherein the solubilizer is present in the composition in an amount of from about 1 to 30% by weight, based on the weight of the composition.

5